

30) Gram-Schmidt

Partial Basis $X = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$

$$\vec{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} - \text{proj}_X \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} - \frac{3}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$4\vec{v}_2 = 4 \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} - 3 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} \quad \text{Partial Basis } X = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} \right\}$$

$$\vec{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \text{proj}_X \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \frac{2}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{(-2)}{12} \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$$

$$6\vec{v}_3 = 6 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - 3 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \\ 4 \\ -2 \end{bmatrix} \quad \text{Orthogonal Basis} = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 4 \\ -2 \end{bmatrix} \right\}$$

31) $A = \lambda_1 \vec{q}_1 \vec{q}_1^T + \lambda_2 \vec{q}_2 \vec{q}_2^T$

where \vec{q}_1, \vec{q}_2 are orthonormal
written as columns.

Section 5.4
eigenvectors,

$$\vec{q}_1 = \frac{1}{\sqrt{10}} \begin{bmatrix} 1 \\ 3 \end{bmatrix} \quad \lambda_1 = 2$$

$$\vec{q}_2 = \frac{1}{\sqrt{10}} \begin{bmatrix} -3 \\ 1 \end{bmatrix} \quad \lambda_2 = -2$$

$$A = \lambda_1 \vec{q}_1 \vec{q}_1^T + \lambda_2 \vec{q}_2 \vec{q}_2^T$$

$$= 2 \frac{1}{\sqrt{10}} \frac{1}{\sqrt{10}} \begin{bmatrix} 1 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 3 \end{bmatrix} - 2 \frac{1}{\sqrt{10}} \frac{1}{\sqrt{10}} \begin{bmatrix} -3 \\ 1 \end{bmatrix} \begin{bmatrix} -3 & 1 \end{bmatrix}$$

$$= \frac{1}{5} \begin{bmatrix} 1 & 3 \\ 3 & 9 \end{bmatrix} - \frac{1}{5} \begin{bmatrix} 9 & -3 \\ -3 & 1 \end{bmatrix}$$

$$= \frac{1}{5} \begin{bmatrix} -8 & 6 \\ 6 & 8 \end{bmatrix}$$

33

$y = a_0 + a_1 x$
 a_0, a_1 are the variables

$$a_0 + a_1 x = y$$

$$1(a_0) + x(a_1) = y$$

$$\begin{bmatrix} 1 & x \\ \vdots & \vdots \\ 1 & 4 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \end{bmatrix} = \begin{bmatrix} y \\ \vdots \\ 7 \end{bmatrix}$$

$$\vec{x}^* = (A^T A)^{-1} A^T \vec{b}$$

$$A^T A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 4 & 10 \\ 10 & 30 \end{bmatrix}$$

$$(A^T A)^{-1} = \frac{1}{20} \begin{bmatrix} 30 & -10 \\ -10 & 4 \end{bmatrix}$$

$$\vec{x}^* = (A^T A)^{-1} A^T \vec{b}$$

$$= \frac{1}{20} \begin{bmatrix} 30 & -10 \\ -10 & 4 \end{bmatrix} \left(\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ 5 \\ 7 \end{bmatrix} \right)$$

$$= \frac{1}{20} \begin{bmatrix} 30 & -10 \\ -10 & 4 \end{bmatrix} \begin{bmatrix} 17 \\ 51 \end{bmatrix}$$

$$= \frac{1}{20} \begin{bmatrix} 0 \\ 34 \end{bmatrix} \leftarrow a_0 = 0$$

$$\leftarrow a_1 = \frac{34}{20} = 1.7$$

$$y = a_0 + a_1 x$$


$$\boxed{y = 1.7x}$$

35

$$|z| = \sqrt{5^2 + 12^2} = 13$$

$$\theta = \tan^{-1} \frac{12}{5} \quad (+\pi?)$$

$$= \tan^{-1} \frac{12}{5}$$



$$z = |z| [\cos \theta + i \sin \theta]$$

$$z = 13 \left[\cos \left(\tan^{-1} \frac{12}{5} \right) + i \sin \left(\tan^{-1} \frac{12}{5} \right) \right]$$

Follow-Up:

Find z^4 .

$$z^4 = 13^4 \left[\cos \left(4 \tan^{-1} \frac{12}{5} \right) + i \sin \left(4 \tan^{-1} \frac{12}{5} \right) \right]$$